

**Vaccination  
Schedule**  
Birth - 18 years

Flu shots are recommended every year starting at 6 months.

Children and teens should stay up-to-date with COVID-19 vaccines.

<p><b><u>Birth</u></b></p> <p>HepB</p>	<p><b><u>12 months</u></b></p> <p>MMR Varicella HepA PCV</p>	<p><b><u>9-10 years</u></b></p> <p>HPV (2 doses) between 9-12 years</p>
<p><b><u>2 months</u></b></p> <p>DTaP-IPV-Hib-HepB PCV Rotavirus (oral)</p>	<p><b><u>15 months</u></b></p> <p>DTaP Hib</p>	<p><b><u>11-12 years</u></b></p> <p>Tdap MenACWY</p>
<p><b><u>4 months</u></b></p> <p>DTaP-IPV-Hib-HepB PCV Rotavirus (oral)</p>	<p><b><u>18-24 months</u></b></p> <p>HepA</p>	<p><b><u>16 years</u></b></p> <p>MenACWY</p>
<p><b><u>6 months</u></b></p> <p>DTaP-IPV-Hib-HepB PCV Rotavirus (oral)</p>	<p><b><u>4 years</u></b></p> <p>MMRV DTaP-IPV</p>	<p><b><u>16-18 years</u></b></p> <p>MenB (2 doses) Discuss with your provider</p>

**Vaccine Key:**

- DTaP: Diphtheria, Tetanus, Pertussis (whooping cough)
- HepA: Hepatitis A
- HepB: Hepatitis B
- Hib: Haemophilus influenzae type b
- HPV: Human Papillomavirus
- IPV: Inactivated Polio Virus
- MenACWY: Meningococcal A, C, W, Y (meningitis)
- MenB: Meningococcal B (meningitis)
- MMR: Measles, Mumps and Rubella
- PCV: Pneumococcal (pneumonia)
- Tdap: Tetanus, Diphtheria, Pertussis (whooping cough)
- Varicella: Chickenpox

**Combination Vaccines (subject to availability):**

- MMRV: Measles Mumps, Rubella and Varicella
- Pediarix: Combination vaccine that includes DTaP-HepB-IPV
- Pentacel: Combination vaccine that includes DTaP-IPV/Hib
- Vaxelis: Combination vaccine that includes DTaP-IPV-Hib-HepB

**More info: [cdc.gov/vaccines](https://www.cdc.gov/vaccines)**